CERTAIN INEQUALITIES INVOLVING THE k-STRUVE FUNCTION

KOTTAKKARAN SOOPPY NISAR, SAIFUL RAHMAN MONDAL, AND JUNESANG CHOI

Abstract. We aim to introduce a k-Struve function and investigate its various properties, including mainly certain inequalities associated this function. One of the inequalities given here is pointed out to be related to the so-called classical Turán type inequality. We also present a differential equation, several recurrence relations, and integral representations for this k-Struve function.

arXiv:1611.09013v1 [math.CA] 28 Nov 2016

1. Introduction and preliminaries

Díaz and Pariguan [4] introduced and investigated the so-called k-gamma function

$$\Gamma_{\mathtt{k}}(x) := \int_0^\infty t^{x-1} e^{-\frac{t^{\mathtt{k}}}{\mathtt{k}}}\, dt \quad \left(\Re(x) > 0;\, \mathtt{k} \in \mathbb{R}^+\right). \tag{1.1}$$

Here and in the following, let $\mathbb{C}$, $\mathbb{R}$, $\mathbb{R}^+$, $\mathbb{N}$, and $\mathbb{Z}^-$ be the sets of complex numbers, real numbers, positive real numbers, positive integers, and negative integers, respectively, and let $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For various properties of the k-gamma function and its applications to generalize other related functions such as k-beta function and k-digamma function, we refer the interested reader, for example, to [4, 6, 7] and the references cited therein.

Nantomah and Prempeh [7] defined the k-digamma function $\Psi_{\mathtt{k}} := \Gamma'_{\mathtt{k}}/\Gamma_{\mathtt{k}}$ whose series representation is given as follows:

$$\Psi_{\mathtt{k}}(t) := \frac{\log \mathtt{k} - \gamma}{\mathtt{k}} - \frac{1}{t} + \sum_{n=1}^{\infty} \frac{t}{n\mathtt{k}(n\mathtt{k}+t)} \quad \left(\mathtt{k} \in \mathbb{R}^+;\, t \in \mathbb{C} \setminus \mathtt{k}\mathbb{Z}^-\right), \tag{1.2}$$

where γ is the Euler-Mascheroni constant (see, e.g., [9, Section 1.2]). A calculation yields

$$\Psi'_{\mathtt{k}}(t) = \sum_{n=0}^{\infty} \frac{1}{(n\mathtt{k}+t)^2} \quad \left(\mathtt{k},\, t \in \mathbb{R}^+\right). \tag{1.3}$$

Clearly, $\Psi_{\mathtt{k}}(t)$ is increasing on $(0, \infty)$.

Turán [10] proved that the Legendre polynomials $P_n(x)$ satisfy the following determinantal inequality

$$\begin{vmatrix} P_n(x) & P_{n+1}(x) \\ P_{n+1}(x) & P_{n+2}(x) \end{vmatrix} \leq 0 \quad (-1 \leq x \leq 1;\, n \in \mathbb{N}_0), \tag{1.4}$$

where the equality occurs only when $x = \pm 1$. Recently, many researchers have applied the above classical inequality (1.4) in various polynomials and functions such as ultraspherical polynomials, Laguerre polynomials, Hermite polynomials, Bessel functions of the first

2010 *Mathematics Subject Classification.* 33C10, 26D07.

Key words and phrases. k-Struve function; k-gamma function; k-beta function; k-digamma function; Turán type inequalities.

kind, modified Bessel functions, and polygamma functions. Karlin and Szegö [5] named such determinants as in (1.4) Turánians.

In this paper, we consider the following k-Struve function (cf. [1, p.496, Entry 12.1.3])

$$\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) := \sum_{r=0}^{\infty} \frac{(-c)^r}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})} \left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1}. \tag{1.5}$$

Then we investigate the k-Struve function (1.5) as follows: We establish certain inequalities involving $\mathtt{S}^{\mathtt{k}}_{\nu,c}$, one of which is shown to be related to the Turán type inequality; we show that the k-Struve function satisfies a second order non-homogeneous differential equation; we present an integral representation and recurrence relations for the k-Struve function.

2. Inequalities

The modified k-Struve function is given as

$$L^{\mathtt{k}}_{\nu}(x) := \mathtt{S}^{\mathtt{k}}_{\nu,-1}(x), \tag{2.1}$$

which is normalized and denoted by $\mathcal{L}^{\mathtt{k}}_{\nu}$ as follows:

$$\mathcal{L}^{\mathtt{k}}_{\nu}(x) = \left(\frac{2}{x}\right)^{\frac{\nu}{\mathtt{k}}} \Gamma_{\mathtt{k}}\left(\nu+\frac{3\mathtt{k}}{2}\right) L^{\mathtt{k}}_{\nu}(x) = \sum_{r=0}^{\infty} f_r(\nu,\mathtt{k})\, x^{2r+1}, \tag{2.2}$$

where

$$f_r(\nu,\mathtt{k}) := \frac{\Gamma_{\mathtt{k}}\left(\nu+\frac{3\mathtt{k}}{2}\right)}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3}{2})\Gamma(r+\frac{3}{2})\, 2^{2r+1}}.$$

Here, we investigate monotonicity and log-convexity involving the $\mathcal{L}^{\mathtt{k}}_{\nu}$. To do this, we recall some known useful properties which are given in the following lemma (see [3]).

Lemma 2.1. *Consider the power series $f(x) = \sum_{\mathtt{k}=0}^{\infty} a_{\mathtt{k}} x^{\mathtt{k}}$ and $g(x) = \sum_{\mathtt{k}=0}^{\infty} b_{\mathtt{k}} x^{\mathtt{k}}$, where $a_{\mathtt{k}} \in \mathbb{R}$ and $b_{\mathtt{k}} \in \mathbb{R}^+$ $(\mathtt{k} \in \mathbb{N}_0)$. Further suppose that both series converge on $|x| < r$. If the sequence $\{a_{\mathtt{k}}/b_{\mathtt{k}}\}_{\mathtt{k}\geq 0}$ is increasing (or decreasing), then the function $x \mapsto f(x)/g(x)$ is also increasing (or decreasing) on $(0, r)$.*

If both f and g are even, or both are odd functions, then the above results will be applicable.

Theorem 2.1. *Let $\mathtt{k} \in \mathbb{R}^+$ be fixed. Then the following statements hold.*

(i) *For $\nu \geq \mu > -3\mathtt{k}/2$, then the function $x \mapsto \mathcal{L}^{\mathtt{k}}_{\mu}(x)/\mathcal{L}^{\mathtt{k}}_{\nu}(x)$ is increasing on $\mathbb{R}$.*

(ii) *The function $\nu \mapsto \mathcal{L}^{\mathtt{k}}_{\nu}(x)$ is decreasing for a fixed $x \in [0,\infty)$ and increasing for a fixed $x \in (-\infty, 0)$. Also, the function $\nu \mapsto \mathcal{L}^{\mathtt{k}}_{\nu}(x)$ is log-convex on $(-3\mathtt{k}/2, \infty)$ for a fixed $x \in \mathbb{R}^+$.*

(iii) *The function $\nu \mapsto L^{\mathtt{k}}_{\nu+\mathtt{k}}(x)/L^{\mathtt{k}}_{\nu}(x)$ is decreasing on $(-3\mathtt{k}/2, \infty)$ for a fixed $x \in \mathbb{R}^+$.*

Proof. To prove (i), recall the series in (2.2). Clearly,

$$\frac{\mathcal{L}_{\nu}^{\mathtt{k}}(x)}{\mathcal{L}_{\mu}^{\mathtt{k}}(x)} = \frac{\sum\limits_{r=0}^{\infty} f_r(\nu,\mathtt{k})\, x^{2r+1}}{\sum\limits_{r=0}^{\infty} f_r(\mu,\mathtt{k})\, x^{2r+1}}.$$

Denote $w_r := f_r(\nu,\mathtt{k})/f_r(\mu,\mathtt{k})$. Then

$$w_r = \frac{\Gamma_{\mathtt{k}}(\nu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(\mu+\frac{3}{2}\mathtt{k})}.$$

Appealing to the relation (3.2), we find

$$\begin{aligned}\frac{w_{r+1}}{w_r} &= \frac{\Gamma_{\mathtt{k}}(\nu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(r\mathtt{k}+\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(\mu+\frac{3}{2}\mathtt{k})}\frac{\Gamma_{\mathtt{k}}(\mu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(\nu+\frac{3}{2}\mathtt{k})}\\ &= \frac{(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})\Gamma(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})}{(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})\Gamma(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})\Gamma_{\mathtt{k}}(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k})} = \frac{r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}}{r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k}} \le 1,\end{aligned}$$

whose last inequality is valid from the condition $\nu \ge \mu > -3\mathtt{k}/2$. Finally, the result (i) follows from Lemma 2.1.

For (ii), since $2\nu > -3\mathtt{k}$, we first observe the coefficients $f_r(\nu,\mathtt{k}) > 0$ for all $r \in \mathbb{N}_0$. Then the logarithmic derivative of $f_r(\nu,\mathtt{k})$ with respect to ν is

$$\frac{f_r'(\nu,\mathtt{k})}{f_r(\nu,\mathtt{k})} = \Psi_{\mathtt{k}}\Big(\nu+\frac{3}{2}\mathtt{k}\Big) - \Psi_{\mathtt{k}}\Big(r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k}\Big) \le 0,$$

whose last inequality follows from (1.2). Since $f_r(\nu,\mathtt{k}) > 0$ ($r \in \mathbb{N}_0$; $2\nu > -3\mathtt{k}$), $f_r{}'(\nu,\mathtt{k}) \le 0$ ($r \in \mathbb{N}_0$; $2\nu > -3\mathtt{k}$). Hence $\nu \mapsto f_r(\nu,\mathtt{k})$ is decreasing on $(-3\mathtt{k}/2,\infty)$. This implies that, for $\mu \ge \nu > -3\mathtt{k}/2$,

$$\sum_{r=0}^{\infty} f_r(\nu,\mathtt{k})x^{2r+1} \ge \sum_{r=0}^{\infty} f_r(\mu,\mathtt{k})x^{2r+1} \quad (x \in [0,\infty))$$

and

$$\sum_{r=0}^{\infty} f_r(\nu,\mathtt{k})x^{2r+1} \le \sum_{r=0}^{\infty} f_r(\mu,\mathtt{k})x^{2r+1} \quad (x \in (-\infty,0)).$$

This proves the first statement of (ii).

In view of (1.3), we have

$$\frac{\partial^2}{\partial\nu^2}\left(\log(f_r(\nu,\mathtt{k}))\right) = \sum_{n=0}^{\infty}\left\{\frac{1}{(n\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})^2} - \frac{1}{(n\mathtt{k}+r\mathtt{k}+\nu+\frac{3}{2}\mathtt{k})^2}\right\} \ge 0$$

for all $\mathtt{k} \in \mathbb{R}^+$ and $2\nu > -3\mathtt{k}$. Therefore $\nu \mapsto f_r(\nu)$ is log-convex on $(-3\mathtt{k}/2,\infty)$. Since a sum of log-convex functions is log-convex, the second statement of (ii) is proved.

For (iii), it is obvious from (i) that

$$\frac{d}{dx}\left(\frac{\mathcal{L}_{\mu}^{\mathtt{k}}(x)}{\mathcal{L}_{\nu}^{\mathtt{k}}(x)}\right) \ge 0, \tag{2.3}$$

for all $x \in \mathbb{R}^+$ and $\nu \geq \mu > -3\mathtt{k}/2$. In view of the relation (2.2), (2.3) is equivalent to

$$\left(x^{-\frac{\mu}{\mathtt{k}}}\mathtt{L}_{\mu}^{\mathtt{k}}(x)\right)'\left(x^{-\frac{\nu}{\mathtt{k}}}\mathtt{L}_{\nu}^{\mathtt{k}}(x)\right) - \left(x^{-\frac{\mu}{\mathtt{k}}}\mathtt{L}_{\mu}^{\mathtt{k}}(x)\right)\left(x^{-\frac{\nu}{\mathtt{k}}}\mathtt{L}_{\nu}^{\mathtt{k}}(x)\right)' \geq 0 \tag{2.4}$$

for all $x \in \mathbb{R}^+$ and $\nu \geq \mu > -3\mathtt{k}/2$.

Considering (2.1) and setting $c = -1$ in (3.4) gives

$$\frac{d}{dx}\left(x^{-\frac{\nu}{\mathtt{k}}}\mathtt{L}_{\nu}^{\mathtt{k}}(x)\right) = \frac{2^{-\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})} + x^{-\frac{\nu}{\mathtt{k}}}\mathtt{L}_{\nu+\mathtt{k}}^{\mathtt{k}}(x). \tag{2.5}$$

Applying (2.5) to the inequality (2.4) and using (2.2), we obtain

$$\begin{aligned} & x^{-\frac{\mu+\nu}{\mathtt{k}}}\left\{\mathtt{L}_{\mu+\mathtt{k}}^{\mathtt{k}}(x)\mathtt{L}_{\nu}^{\mathtt{k}}(x) - \mathtt{L}_{\nu+\mathtt{k}}^{\mathtt{k}}(x)\mathtt{L}_{\mu}^{\mathtt{k}}(x)\right\} \\ & \geq \frac{2^{-\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})}x^{-\frac{\mu}{\mathtt{k}}}\mathtt{L}_{\mu}^{\mathtt{k}}(x) - \frac{2^{-\frac{\mu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\mu+\frac{3\mathtt{k}}{2})}x^{-\frac{\nu}{\mathtt{k}}}\mathtt{L}_{\nu}^{\mathtt{k}}(x) \\ & = \frac{2^{-\frac{\mu+\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\mu+\frac{3\mathtt{k}}{2})\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})}\left(\mathcal{L}_{\mu}^{\mathtt{k}}(x) - \mathcal{L}_{\nu}^{\mathtt{k}}(x)\right) \geq 0 \end{aligned} \tag{2.6}$$

for all $x \in \mathbb{R}^+$ and $\nu \geq \mu > -3\mathtt{k}/2$. Here, the last inequality in (2.6) follows from the first statement of (ii). Also, we find from (2.6) that

$$\frac{L_{\mu+\mathtt{k}}^{\mathtt{k}}(x)}{L_{\mu}^{\mathtt{k}}(x)} - \frac{L_{\nu+\mathtt{k}}^{\mathtt{k}}(x)}{L_{\nu}^{\mathtt{k}}(x)} \geq 0$$

for all $x \in \mathbb{R}^+$ and $\nu \geq \mu > -3\mathtt{k}/2$. This proves (iii). □

Remark 2.1. One of the most significant consequences of Theorem 2.1 is the Turán-type inequality for the function $\mathcal{L}_{\nu}^{\mathtt{k}}$. The log-convexity of $\mathcal{L}_{\nu}^{\mathtt{k}}$ (the last statement of (ii) in Theorem 2.1) implies

$$\mathcal{L}_{\alpha\nu_1+(1-\alpha)\nu_2}^{\mathtt{k}}(x) \leq \left(\mathcal{L}_{\nu_1}^{\mathtt{k}}\right)^{\alpha}(x)\left(\mathcal{L}_{\nu_2}^{\mathtt{k}}\right)^{1-\alpha}(x) \tag{2.7}$$

$$\left(\alpha \in [0,1];\ x,\, \mathtt{k} \in \mathbb{R}^+;\ \nu_1,\, \nu_2 \in (-3\mathtt{k}/2, \infty)\right).$$

Choosing $\alpha = 1/2$ and setting $\nu_1 = \nu - \mathtt{a}$ and $\nu_2 = \nu + \mathtt{a}$ for some $\mathtt{a} \in \mathbb{R}$ in (2.7) yields the following reversed Turán type inequality (cf. (1.4))

$$\left(\mathcal{L}_{\nu}^{\mathtt{k}}(x)\right)^2 - \mathcal{L}_{\nu-\mathtt{a}}^{\mathtt{k}}(x)\mathcal{L}_{\nu+\mathtt{a}}^{\mathtt{k}}(x) \leq 0 \tag{2.8}$$

$$\left(x,\, \mathtt{k} \in \mathbb{R}^+;\ \mathtt{a} \in \mathbb{R},\, \nu \in (|\mathtt{a}| - 3\mathtt{k}/2, \infty)\right).$$

3. Formulae for the k-Struve function

Here, we present a differential equation, and recurrence relations regarding the k-Struve function $\mathtt{S}_{\nu,c}^{\mathtt{k}}$ (1.5).

Proposition 3.1. *Let $\mathtt{k} \in \mathbb{R}^+$ and $2\nu > -3\mathtt{k}$. Then the k-Struve function $\mathtt{S}_{\nu,c}^{\mathtt{k}}$ (1.5) satisfies the following second-order non-homogeneous differential equation*

$$x^2\frac{d^2y}{dx^2} + x\frac{dy}{dx} + \frac{1}{\mathtt{k}^2}\left(c\,\mathtt{k}\,x^2 - \nu^2\right)y = \frac{4\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}+1}}{\mathtt{k}\Gamma_{\mathtt{k}}(\nu+\frac{\mathtt{k}}{2})\Gamma(\frac{1}{2})}. \tag{3.1}$$

Proof. By using the k-Struve function $\mathtt{S}^{\mathtt{k}}_{\nu,c}$ (1.5) and the functional relation

$$\Gamma_{\mathtt{k}}(x+\mathtt{k}) = x\,\Gamma_{\mathtt{k}}(x), \tag{3.2}$$

we find

$$\begin{aligned}
&x^2\frac{d^2}{dx^2}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) + x\frac{d}{dx}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\\
&= \sum_{r=0}^{\infty}\frac{(-c)^r\left(2r+\frac{\nu}{\mathtt{k}}+1\right)^2}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1}\\
&= \sum_{r=0}^{\infty}\frac{(-c)^r(2r+1)\left(2r+\frac{2\nu}{\mathtt{k}}+1\right)}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1} + \frac{\nu^2}{\mathtt{k}^2}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\\
&= \frac{4}{\mathtt{k}}\sum_{r=0}^{\infty}\frac{(-c)^r\left(r+\frac{1}{2}\right)\left(r\mathtt{k}+\nu+\frac{\mathtt{k}}{2}\right)}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1} + \frac{\nu^2}{\mathtt{k}^2}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\\
&= \frac{4}{\mathtt{k}}\frac{\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}+1}}{\Gamma_{\mathtt{k}}(\nu+\frac{\mathtt{k}}{2})\Gamma(\frac{1}{2})} + \frac{4}{\mathtt{k}}\sum_{r=1}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{\mathtt{k}}{2})\Gamma(r+\frac{1}{2})}\left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1} + \frac{\nu^2}{\mathtt{k}^2}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\\
&= \frac{4}{\mathtt{k}}\frac{\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}+1}}{\Gamma_{\mathtt{k}}(\nu+\frac{\mathtt{k}}{2})\Gamma(\frac{1}{2})} - \frac{cx^2}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) + \frac{\nu^2}{\mathtt{k}^2}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x).
\end{aligned}$$

This shows that $y = \mathtt{S}^{\mathtt{k}}_{\nu,c}(x)$ the differential equation (3.1). □

Theorem 3.1. *Let $\mathtt{k} \in \mathbb{R}^+$ and $2\nu > -3\mathtt{k}$. Then the following recurrence relations hold true:*

$$\frac{d}{dx}\left(x^{\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right) = \frac{1}{\mathtt{k}}x^{\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu-\mathtt{k}}(x); \tag{3.3}$$

$$\frac{d}{dx}\left(x^{-\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right) = \frac{2^{-\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})} - c\,x^{-\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu+\mathtt{k},c}(x); \tag{3.4}$$

$$\frac{1}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu-\mathtt{k}}(x) - c\,\mathtt{S}^{\mathtt{k}}_{\nu+\mathtt{k},c}(x) = 2\frac{d}{dx}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) - \frac{(x/2)^{\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})}; \tag{3.5}$$

$$\frac{1}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu-\mathtt{k}}(x) + c\,\mathtt{S}^{\mathtt{k}}_{\nu+\mathtt{k},c}(x) = \frac{2\nu}{x\,\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) + \frac{(x/2)^{\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})}. \tag{3.6}$$

Proof. From (3.1) we have

$$x^{\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) = \sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\frac{x^{2r+\frac{2\nu}{\mathtt{k}}+1}}{2^{2r+\frac{\nu}{\mathtt{k}}+1}},$$

which, upon differentiating with respect to x and using the relation (3.2), yields

$$\begin{aligned}
\frac{d}{dx}\left(x^{\frac{\nu}{\mathtt{k}}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right) &= \sum_{r=0}^{\infty}\frac{(-c)^r(2r+\frac{2\nu}{\mathtt{k}}+1)}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\frac{x^{2r+\frac{2\nu}{\mathtt{k}}}}{2^{2r+\frac{\nu}{\mathtt{k}}+1}}\\
&= \frac{x^{\frac{\nu}{\mathtt{k}}}}{\mathtt{k}}\sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{\mathtt{k}}{2})\Gamma(r+\frac{3}{2})}\left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}} = \frac{1}{\mathtt{k}}\,x^{\frac{\nu}{\mathtt{k}}}\,\mathtt{S}^{\mathtt{k}}_{\nu-\mathtt{k},c}(x).
\end{aligned}$$

This prove (3.3). We can establish the result (3.4) by a similar argument as in the proof of (3.3). We omit the details.

Similarly, from (1.5), we obtain

$$x\frac{d}{dx}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)+\frac{\nu}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)=\frac{x}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu-\mathtt{k}}(x) \tag{3.7}$$

and

$$x\frac{d}{dx}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)-\frac{\nu}{\mathtt{k}}\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)=\frac{x(x/2)^{\frac{\nu}{\mathtt{k}}}}{\sqrt{\pi}\,\Gamma_{\mathtt{k}}(\nu+\frac{3\mathtt{k}}{2})}-c\,x\mathtt{S}^{\mathtt{k}}_{\nu+\mathtt{k},c}(x). \tag{3.8}$$

Adding and subtracting each side of (3.7) and (3.8) yields, respectively, the results (3.5) and (3.6).

□

4. Integral representations

Here, we present two integral representations for the function $\mathtt{S}^{\mathtt{k}}_{\nu,c}$.

Theorem 4.1. *Let* $\mathtt{k}\in\mathbb{R}^+$, $\Re(\nu)>-\frac{\mathtt{k}}{2}$, *and* $\alpha\in\mathbb{R}\setminus\{0\}$. *Then*

$$\mathtt{S}^{\mathtt{k}}_{\nu,\alpha^2}(x)=\frac{2\sqrt{\mathtt{k}}}{\alpha^2\sqrt{\pi}\Gamma_{\mathtt{k}}\left(\nu+\frac{\mathtt{k}}{2}\right)}\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}}\int_0^1(1-t^2)^{\frac{\nu}{\mathtt{k}}-\frac{1}{2}}\sin\left(\frac{\alpha xt}{\sqrt{\mathtt{k}}}\right)dt. \tag{4.1}$$

and

$$\mathtt{S}^{\mathtt{k}}_{\nu,-\alpha^2}(x)=\frac{2\sqrt{\mathtt{k}}}{\sqrt{\pi}\Gamma_{\mathtt{k}}\left(\nu+\frac{\mathtt{k}}{2}\right)}\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}}\int_0^1(1-t^2)^{\frac{\nu}{\mathtt{k}}-\frac{1}{2}}\sinh\left(\frac{\alpha xt}{\sqrt{\mathtt{k}}}\right)dt. \tag{4.2}$$

In particular, we have

$$1-\cos\left(\frac{\alpha x}{\sqrt{\mathtt{k}}}\right)=\frac{\alpha}{\mathtt{k}}\sqrt{\frac{\pi x}{2}}\mathtt{S}^{\mathtt{k}}_{\frac{1}{2},\alpha^2}(x) \tag{4.3}$$

and

$$\cosh\left(\frac{\alpha x}{\sqrt{\mathtt{k}}}\right)-1=\frac{\alpha}{\mathtt{k}}\sqrt{\frac{\pi x}{2}}\mathtt{S}^{\mathtt{k}}_{\frac{\nu}{\mathtt{k}},-\alpha^2}(x). \tag{4.4}$$

Proof. We begin by recalling the $\mathtt{k}$-beta function (see [4])

$$\mathtt{B}_{\mathtt{k}}(x,y)=\frac{\Gamma_{\mathtt{k}}(x)\Gamma_{\mathtt{k}}(y)}{\Gamma_{\mathtt{k}}(x+y)}=\frac{1}{\mathtt{k}}\int_0^1 t^{\frac{x}{\mathtt{k}}-1}(1-t)^{\frac{y}{\mathtt{k}}-1}\,dt \tag{4.5}$$

$$\left(\mathtt{k}\in\mathbb{R}^+;\,\min\{\Re(x),\,\Re(y)\}>0\right).$$

Replacing t by t^2 on the right-hand-sided integral in (4.5), we obtain

$$\mathtt{B}_{\mathtt{k}}(x,y)=\frac{2}{\mathtt{k}}\int_0^1 t^{\frac{2x}{\mathtt{k}}-1}(1-t^2)^{\frac{y}{\mathtt{k}}-1}\,dt. \tag{4.6}$$

Setting $x=(r+1)\mathtt{k}$ and $y=\nu+\mathtt{k}/2$ in (4.6) gives

$$\frac{1}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\mathtt{k})}=\frac{2}{\Gamma_{\mathtt{k}}\left((r+1)\,\mathtt{k}\right)\Gamma_{\mathtt{k}}\left(\nu+\frac{\mathtt{k}}{2}\right)}\int_0^1 t^{2r}(1-t^2)^{\frac{\nu}{\mathtt{k}}-\frac{1}{2}}\,dt. \tag{4.7}$$

Applying the known identity

$$\Gamma_{\mathtt{k}}(\mathtt{k}x)=\mathtt{k}^{x-1}\Gamma(x) \tag{4.8}$$

and the Legendre duplication formula (see [1, 2, 9])

$$\Gamma(z)\Gamma\left(z+\tfrac{1}{2}\right)=2^{1-2z}\,\sqrt{\pi}\,\Gamma(2z) \tag{4.9}$$

to the function $\mathtt{S}^{\mathtt{k}}_{\nu,c}$ with (4.7), we get

$$\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)=\frac{2\sqrt{\mathtt{k}}}{\sqrt{\pi}\Gamma_{\mathtt{k}}\left(\nu+\frac{\mathtt{k}}{2}\right)}\left(\frac{x}{2}\right)^{\frac{\nu}{\mathtt{k}}}\int_0^1(1-t^2)^{\frac{\nu}{\mathtt{k}}-\frac{1}{2}}\sum_{r=0}^{\infty}\frac{(-c)^r}{(2r+1)!}\left(\frac{xt}{\sqrt{\mathtt{k}}}\right)^{2r+1}dt. \tag{4.10}$$

Finally, setting $c=\pm\alpha^2$ ($\alpha\in\mathbb{R}\setminus\{0\}$) in (4.10) yields, respectively, the desired results (4.1) and (4.2). Further, setting $\nu=\mathtt{k}/2$ in (4.1) and (4.2) yields, respectively, the desired results (4.3) and (4.4).

□

References

[1] M. Abramowitz and I.A. Stegun (Editors), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Tenth Printing, National Bureau of Standards, Applied Mathematics Series **55**, National Bureau of Standards, Washington, D.C., 1972; Reprinted by Dover Publications, New York, 1965 (see also [8]).

[2] G.E. Andrews, R. Askey and R. Roy, *Special Functions*, Encyclopedia of Mathematics and Its Applications, Vol. **71**, Cambridge University Press, Cambridge, London and New York, 1999.

[3] M. Biernacki and J. Krzyż, On the monotonity of certain functionals in the theory of analytic functions, *Ann. Univ. Mariae Curie-Sk lodowska. Sect. A.* **9** (1957), 135-147.

[4] R. Díaz and E. Pariguan, On hypergeometric functions and k-Pochhammer symbol, *Divulg. Mat.* **15**(2) (2007), 179-192.

[5] S. Karlin and G. Szegö, On certain determinants whose elements are orthogonal polynomials, *J. Analyse Math.* **8** (1960/1961), 1-157.

bibitemMi D.S. Mitrinović, *Analytic Inequalities*, Springer, New York, 1970.

[6] S. Mubeen, M. Naz and G. Rahman, A note on k-hypergemetric differential equations, *J. Inequal. Spec. Funct.* **4**(3) (2013), 8-43.

[7] K. Nantomah and E. Prempeh, Some inequalities for the k-Digamma function, *Mathematica Aeterna* **4**(5) (2014), 521-525.

[8] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark (Editors), *NIST Handbook of Mathematical Functions* [With 1 CD-ROM (Windows, Macintosh and UNIX)], U. S. Department of Commerce, National Institute of Standards and Technology, Washington, D. C., 2010; Cambridge University Press, Cambridge, London and New York, 2010 (see also [1]).

[9] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

[10] P. Turán, On the zeros of the polynomials of Legendre, *Časopis Pěst. Mat. Fys.* **75** (1950), 113-122.

Department of Mathematics, College of Arts and Science-Wadi Al dawser, 11991,, Prince Sattam bin Abdulaziz University, Saudi Arabia
E-mail address: ksnisar1@gmail.com, n.sooppy@psau.edu.sa

Department of Mathematics, King Faisal University, Al Ahsa 31982, Saudi Arabia
E-mail address: smondal@kfu.edu.sa

Department of Mathematics, Dongguk University, Gyeongju 38066, Republic of Korea
E-mail address: junesang@mail.dongguk.ac.kr